

PhD Thesis

THE ANTARCTIC-MAGELLAN CONNECTION: DIVERSITY AND BIOGEOGRAPHY OF INTERSTITIAL TURBELLARIANS IN THE SCOTIA ARC

M.Sc. Odile Volonterio

Sección Zoología de Invertebrados, Facultad de Ciencias, Universidad de la República

E-mail: odile@fcien.edu.uy

INTRODUCTION:

The study of systematics and distribution of Antarctic organisms is very important, not only for a few specialists, but also for all biologists involved in biogeography (Dell, 1972, *sensu* Karling, 1977). Despite the relevance of the relation between the Antarctic and Magellanic biotas, its study has proven difficult because of substantial problems that make the biogeographical evaluations rest on incomplete data (Arntz, 1999; Convey, 2007). Main problems include (i) the lack of specialists, because of which in many works the determinations do not reach species level; (ii) the lack of studies in the Scotia Arc region, that results in that many works compare the Magellanic Region with the Weddell Sea, because the latter has been better investigated, and (iii) the low biodiversity in the Antarctic (Clarke *et al.*, 2007). Furthermore, in recent years it has become clear that studies based on different groups of organisms can yield contradictory results (Griffiths *et al.*, 2009), especially when comparing distributions of taxa with dispersive phases (for example, see Brandt *et al.*, 1999; Liuzzi and Zelaya, 2004; Schrödl, 1999, and Zelaya, 2005 for mollusk distributions).

The study of turbellarians can overcome the latter difficulty. Because of their adaptations to the interstitial environment, these animals have a practically null dispersal potential, which makes vicariance processes the most likely to explain their current distribution. There are three examples of the use of these organisms in large-scale biogeographical studies, and all three have led to consistent results: (i) the work of Sterrer (1973), where trans-allopatric sister species of many interstitial phyla, including turbellarians, were found on both sides of the Atlantic; (ii) two complementary works by Sluys (1989, 1995), who found evidence of trans-Pacific tracks using marine and terrestrial planarians, and (iii) two complementary works by Ax and Armonies (1987, 1990), who found evidence of an ancient boreal circumpolar distribution of brackish water turbellarians.

This study will provide significant data on the diversity and distribution of a group of organisms that has been overlooked in most studies of the benthic fauna of the Antarctic Region, and it will also attempt to provide clues for the evolution of the Antarctic biota in a region of key importance for biogeographical studies. With the Antarctic Peninsula being one of the most rapidly changing environments in the world, it is imperative to undertake these kind of analysis as soon as possible, in order to get reliable data before both human impact and climatic change alter the biogeographic signals provided by the biota.

OBJECTIVES:

The general objective of this proposal is to carry out a Historical Biogeography study to reconstruct the history of the biota of marine interstitial turbellarians in the Scotia Arc. In order to do so, four specific objectives have been defined: (i) to describe the biota of interstitial turbellarians in the Scotia Arc area, based on sampling in the Antarctic Peninsula, King George Island, Tierra del Fuego and the Magellan Strait, and on reports from the South Georgia, South Orkney and the Falkland/Malvinas Islands; (ii) to make a panbiogeographic analysis using this fauna, to test the processes of vicariance suggested for the area by the available geological information; (iii) to carry out an analysis of predictive modelling of species distribution to complement the biogeographical analysis, and (iv) to evaluate the results in the context of available literature.

RELEVANCE:

This proposal falls within the goals of the Evolution and Biodiversity in the Antarctic (EBA) Scientific Research Programme.

METHODOLOGY:

Sampling. The Uruguayan Antarctic Institute will facilitate sampling in the Antarctic Peninsula, King George Island, Tierra del Fuego and the Magellan Strait. Samples will be processed mainly in the Uruguayan Base Científica Antártica Artigas (King George Island), and also in the Uruguayan Estación Científica Antártica T/N Ruperto Elichiribehety (Hope Bay, Antarctic Peninsula).

Taxonomical study. Species will be identified using standard techniques for the study of microinvertebrates, including the observation of live animals, and morphometrical and histological analyses.

Biogeographical analysis. A panbiogeographical analysis will be carried out under a vicariance paradigm. Individual tracks will be drawn at species and genus level. Orientation of tracks will be performed with the baseline and main massing criteria. Generalized tracks will be obtained, and nodes identified. A complementary analysis of predictive modelling of species distribution will be carried out with informatic tools.

Interpretation of results. The results will be analyzed in the context of (i) the distribution of species and genera of interstitial turbellarians at a global level, (ii) the proposed evolution of the Scotia Arc area, and (iii) other biogeographical analyses performed in the area.

EXPECTED RESULTS:

Internal deliverables include a Collection of specimens identified at the species level, which will be deposited in the Helminthological Collection of the Faculty of Sciences, Montevideo, Uruguay, a taxonomic database of anatomy and morphometry of the species, and internal reports.

External deliverables include a set of specimens that will be deposited in an external Collection, a list of the species of marine interstitial turbellarians in different localities along the Scotia Arc, taxonomic publications describing taxa, a report and a publication of the results of the biogeographical study, and recommendations for the delimitation of protected areas.

REFERENCES CITED:

- Arntz WE. 1999. Magellan-Antarctic: Ecosystems that drifted apart. Summary Review. *Scientia Marina* 63 (Supl. I): 503-511.
- Ax P and W Armonies. 1987. Amphiatlantic identities in the composition of the boreal brackish water community of Platyhelminthes. *Microfauna Marina* 3: 7-80.
- Ax P and W Armonies. 1990. Brackish water Platyhelminthes from Alaska as evidence for the existence of a boreal brackish water community with circumpolar distribution. *Microfauna Marina* 6: 7-109.
- Brandt A, Linse K and U Mühlenhardt-Siegel. 1999. Biogeography of Crustacea and Mollusca of the Subantarctic and Antarctic regions. *Scientia Marina* 63 (Supl. 1): 383-389.
- Clarke A, Griffiths HJ, Linse K, Barnes DKA y JA Crame. 2007. How well do we know the Antarctic marine fauna? A preliminary study of macroecological and biogeographical patterns in Southern Ocean gastropod and bivalve mollusks. *Diversity and Distributions* 13: 620-632.
- Convey P. 2007. Influences on and origins of terrestrial biodiversity of the sub-Antarctic islands. *Papers and Proceedings of the Royal Society of Tasmania* 141: 83-93.
- EBA, 2009. Evolution and Biodiversity in the Antarctic. Accessed 12/05/2009. <http://www.eba.aq/>
- Griffiths HJ, Barnes DKA y K Linse. 2009. Towards a generalized biogeography of the Southern Ocean benthos. *Journal of Biogeography* 36: 162-177.
- Karling TG. 1977. Taxonomy, Phylogeny and Biogeography of the genus *Austrorhynchus* Karling (Turbellaria, Polycystididae). En: *The Meiofauna species in time and space. Proceedings of a Workshop Symposium. Bermuda Biological Station, 1975*. Sterrer, W y P Ax (Eds.). *Mikrofauna des Meeresbodens* 61: 153-165.
- Liuzzi MG and DG Zelaya. 2004. El rol de las islas Georgias del Sur en la distribución de los poliplacóforos del Océano Sur. Resumen del 5° Simposio Argentino y 1° Latinoamericano sobre Investigaciones Antárticas. Accessed 27/04/2009: <http://www.dna.gov.ar/CIENCIA/SANTAR04/CD/PDF/202BP.PDF>
- Schrödl M. 1999. Zoogeographic relationships of Magellan Nudibranchia (Mollusca: Opisthobranchia) with particular reference to species from adjacent regions. *Scientia Marina* 63 (Supl. 1): 409-416.
- Sluys R. 1989. A monograph of the marine triclads. Balkema, Rotterdam, Brookfield, 463 pp.
- Sluys R. 1995. Platyhelminths as paleogeographical indicators. *Hydrobiologia* 305: 49-53.
- Sterrer W. 1973. Plate tectonics as a mechanism for dispersal and speciation in interstitial sand fauna. *Netherlands Journal of Sea Research* 7: 200-222.
- Zelaya DG. 2005. The bivalves from the Scotia Arc islands: Species richness and faunistic affinities. *Scientia Marina* 69 (Supl. 2): 113-122.